


METÓDY RIEŠENIA FYZIKÁLNYCH ÚLOH 2 zima24 – Príklady 4


Cvičenie 13.11.2024

Príklad 1

A block of mass M_h slides without friction on a horizontal table. It is connected by a massless rope passing over a massless frictionless pulley to a second hanging mass M_v pulled downward by gravity. A sphere of mass m and radius R , initially at rest, rolls without sliding on the top surface of the first block. Find the resulting acceleration of the mass M_v and the center of mass of the sphere.




Two uniform cylinders spin independently about their axes (the axes are parallel to each other). The first has radius r_1 and mass m_1 , the other has radius r_2 and mass m_2 . Initially they rotate in the same sense of rotation with angular speeds ω_1 and ω_2 respectively. They are then brought together so that they touch. After the steady state is achieved, what is the final angular velocity of cylinder 1, ω'_1 ?



Príklad 2


Find the heat capacity of the system consisting of a container that has two compartments separated by a piston. To the right of the piston is a gas of monoatomic molecules (gas parameters p_0, T_0, V_0), and to the left is vacuum. The piston is held by a spring. If the gas is removed, the piston touches the right wall and the spring is relaxed.



Heat capacities of the materials composing spring, piston and container walls can be neglected.

Príklad 3

Two masses m separated by a distance l are given initial velocities v_0 as shown in the diagram. The masses interact only through universal gravitation.



- Under what conditions will the masses eventually collide?
- Under what conditions will the masses follow circular orbits of diameter l ?
- Under what conditions will the masses follow closed orbits?
- What is the minimum distance achieved between the masses along their path?